

# Chemotherapy Dose Calculation and Administration in Exotic Animal Species

Kevin A. Hahn, DVM, PhD, Dip. ACVIM (Oncology)

## Abstract

There is little information in the literature regarding the use of chemotherapy to treat cancer in exotic animals. This article provides a historical perspective on the use and utility of the body surface area scheme for dosing chemotherapy in animals. Normogram-based recommendations for arriving at a proper chemotherapy dose for the management of cancer in exotic animals are made. It is realistic to offer treatment for many neoplastic diseases in exotic animal species, as long as the limitations are realized and informed consent of the owner is obtained. Copyright 2005 Elsevier Inc. All rights reserved.

**Key words:** Ferrets; avians; exotics; chemotherapy; dosing; body surface area

Administration of a drug to a patient carries with it the implicit assumption that the drug will do something to that patient. There are 2 possible clinical outcomes related to drug delivery. The first of these is therapeutic and desirable, whereas the other is toxic and not desirable. The obvious goal in treating a patient with a drug is to maximize the likelihood of producing a therapeutic response while minimizing the likelihood of producing unacceptable toxicity. The dosing variables available to achieve this goal include the amount of drug delivered and the interval or frequency at which the drug can be given. Under ideal circumstances, these variables are based on sound knowledge of the relationship between the dose, or concentration, of the agent; the likelihood of therapeutic and toxic consequences resulting from its delivery; and the duration of drug effects. Unfortunately, such precise information is lacking for most antineoplastic chemotherapeutic agents. Moreover, it is obvious that not all patients are the same. As a result of genotypic and phenotypic differences, the same dose of drug will produce a range of concentration-versus-time profiles in any given group of patients, with the resulting range of therapeutic and toxic responses corresponding to that pharmacokinetic variability.<sup>1-6</sup>

For drugs that produce therapeutic effects at doses far less than those that cause toxicity, the incentive for precise dosing is far less than for drugs with a narrower therapeutic index. The narrow therapeutic index of most antineoplastic agents has provided great impetus to deliver doses as precisely as possible. One of the practices embedded in dosing of antitumor drugs is dosing by body surface area (BSA), most commonly milligrams per square meter ( $\text{mg}/\text{m}^2$ ). BSA is equivalent to the two-dimensional surface area of the skin. It is difficult to measure, and therefore commonly estimated on the basis of formulas that use body weight and body length in the calculation. The most commonly used formula was published by Du Bois and Du Bois in 1916.<sup>7</sup> Obviously, the objective at that time was not to develop a

---

From Gulf Coast Veterinary Specialists, 1111 West Loop South, Suite 150, Houston, TX 77027 USA.

Address correspondence to: Kevin A. Hahn, DVM, PhD, Dip. ACVIM (Oncology), Gulf Coast Veterinary Specialists, 1111 West Loop South, Suite 150, Houston, TX 77027 USA. E-mail: drhahn@gcvcs.com

© 2005 Elsevier Inc. All rights reserved.

1055-937X/05/1403-\$30.00

doi:10.1053/j.saep.2005.06.004

**Table 1. Five formulas for calculating body surface area (BSA), ranked according to the root-mean-squared-error method of prediction by Wang et al.<sup>3</sup>**

Author	BSA formula
Boyd††	$BSA (m^2) = Wt(kg)^{0.4838} \cdot Ht(cm)^{0.3} \cdot 0.017827$
Gehan and George†	$BSA (m^2) = Wt(kg)^{0.51456} \cdot Ht(cm)^{0.42246} \cdot 0.02350$
Mosteller‡	$BSA (m^2) = [Ht(cm) \cdot Wt(kg)/3600]^{1/2}$ or $BSA (m^2) = [Ht(in) \cdot Wt(lbs)/3131]^{1/2}$
Haycock§	$BSA (m^2) = Wt(kg)^{0.5378} \cdot Ht(cm)^{0.3964} \cdot 0.024265$
Du Bois and Du Bois	$BSA (m^2) = Wt(kg)^{0.425} \cdot Ht(cm)^{0.725} \cdot 0.007184$

††This formula is based on 197 observations.<sup>27</sup>  
†This formula is based on direct measurements of 401 individuals.<sup>23</sup>  
‡This formula is a simple modification of the equation by Gehan and George.<sup>23,28</sup>  
§This formula is based on measurements of 81 individuals, ranging from premature infants to adults.<sup>29</sup>  
||This formula is based on measurements of 9 individuals, one of whom was a child.<sup>7</sup>  
\*Wang Y, Moss J, and Thisted R: Predictors of body surface area. *J Clin Anesth* 4:4-10, 1992

formula to dose anticancer agents, because no such drugs yet existed; Du Bois and Du Bois were working on "clinical calorimetry" (now known as basal metabolic rate). The BSA of mammals correlates with basal metabolic rate. As may be expected in warm-blooded animals, BSA is also proportional to blood volume. But, as Baker and coworkers point out, BSA is not well correlated with glomerular filtration rate.<sup>8,9</sup> BSA is also not associated with liver function.<sup>10</sup> The practice of using BSA in scaling drug doses began with Freireich and coworkers, who quantitatively compared toxicity of anticancer agents in the mouse, rat, hamster, dog, monkey, and human.<sup>11</sup> This introduced the use of BSA in scaling a dose from a mouse or other laboratory animal to an initial starting dose for a phase I study in humans.<sup>12</sup> BSA-based dosing eventually found its way to become the requirement for Food and Drug Administration-approved labeling. Subsequent generations of oncologists also viewed BSA-based dosing as the standard for safe and effective administration of cytotoxic chemotherapy.

What may have been lost in this nearly 40-year-old practice is that the difference in size between mouse and man, or dog and man, is far greater than the variability in size among patients.<sup>10,13-15</sup> As with many practices ingrained in the practice of medicine, mg/m<sup>2</sup> dosing of antitumor drugs has become accepted without questioning the soundness or validity of its underlying assumptions, despite the appearance and availability of newer technologies and theories that would make testing of such assumptions possible.<sup>1-6,16</sup> As analytic chemical instrumentation, pharmacokinetic modeling, and increasingly sophisticated means of as-

sessing molecular and clinical outcomes of drug therapy have been developed, there has been an increasing call to test the validity of the assumptions behind mg/m<sup>2</sup> dosing of antitumor drugs to human and animal species.<sup>3,4,6,16</sup>

## Chemotherapy Dosing in Exotics

There is little information in the literature regarding the use of chemotherapy to treat cancer in exotic animals.<sup>17</sup> Much of the literature that does exist relies on extrapolation from the human literature or the treatment of dogs and cats, is presented as case reports, and includes inadequate follow-up information. Many articles describe the toxicity of chemotherapeutic agents in rabbits, rats, and mice, but the drugs were often given to healthy animals and often at doses that were not therapeutic.

Nevertheless, it is still possible to offer treatment for many neoplastic diseases in exotic animal species, as long as the limitations are realized and informed consent of the owner is obtained.<sup>17,18</sup> It is important to understand the basic mechanisms of action, potential toxicities, and personnel protection issues before attempting to treat any patient of any species with chemotherapy. Clients should be made aware that there are no currently approved chemotherapy agents for use in exotic animal species, and that these drugs are being used in an experimental manner for compassionate purposes. This means informing clients that in most cases dosing information is limited, and all potential toxicities are yet to be elucidated. This also means that limited informa-

**Table 2. Body surface area dosing in dogs and cats using a modification of the Du Bois and Du Bois formula ( $m^2 = 10.0 \times (\text{weight in grams})^{2/3} / 10000$ ).<sup>1-7,16</sup>**

kg	lb	m <sup>2</sup>	kg	lb	m <sup>2</sup>
0.50	1.1	0.06	33	72.6	1.03
1	2.2	0.10	34	74.8	1.05
2	4.4	0.15	35	77.0	1.07
3	6.6	0.20	36	79.2	1.09
4	8.8	0.25	37	81.4	1.11
5	11.0	0.29	38	83.6	1.13
6	13.2	0.33	39	85.8	1.15
7	15.4	0.36	40	88.0	1.17
8	17.6	0.40	41	90.2	1.19
9	19.8	0.43	42	92.4	1.21
10	22.0	0.46	43	94.6	1.23
11	24.2	0.49	44	96.8	1.25
12	26.4	0.52	45	99.0	1.26
13	28.6	0.55	46	101.2	1.28
14	30.8	0.58	47	103.4	1.30
15	33.0	0.60	48	105.6	1.32
16	35.2	0.63	49	107.8	1.34
17	37.4	0.66	50	110.0	1.36
18	39.6	0.69	52	112.2	1.41
19	41.8	0.71	54	114.4	1.44
20	44.0	0.74	56	116.6	1.48
21	46.2	0.76	58	118.8	1.51
22	48.4	0.78	60	121.0	1.55
23	50.6	0.81	62	123.2	1.58
24	52.8	0.83	64	125.4	1.62
25	55.0	0.85	66	127.6	1.65
26	57.2	0.88	68	129.8	1.68
27	59.4	0.90	70	132.0	1.72
28	61.6	0.92	72	134.2	1.75
29	63.8	0.94	74	136.4	1.78
30	66.0	0.96	76	138.6	1.81
31	68.2	0.99	78	140.8	1.84
32	70.4	1.01	80	143.0	1.88

tion is available with regard to prognosis in most cases.

Only doxorubicin and the platinum-containing chemotherapeutic agents (carboplatin, cisplatin) have been studied in veterinary medicine to determine if dosing by BSA would reduce interpatient variability and toxicoses, and none of these drugs was found to have significant relationships between their pharmacokinetics and BSA.<sup>1-6,16,19,20</sup> Results to date suggest that veterinary BSA estimates may be inaccurate, because the values for

**Table 3. Representative surface area to weight ratios (Km) for various species.<sup>11</sup>**

Species	Body weight (kg)	Surface area (m <sup>2</sup> )	Km factor
Mouse	0.02	0.0066	3.0
Rat	0.15	0.025	5.9
Monkey	3.0	0.24	12
Dog	8.0	0.4	20
Human, child	20	0.8	25
Human, adult	60	1.6	37

*Example: To express a mg/kg dose in any given species as the equivalent mg/m<sup>2</sup> dose, multiply the dose by the appropriate Km factor. In the adult human, 100 mg/kg is equivalent to 100 mg/kg  $\times$  37 kg/m<sup>2</sup> = 3700 mg/m<sup>2</sup>.*

the constant (K) and exponent (a) in the formula ( $BSA = K \cdot W^a$ ) are incorrect or because a linear parameter such as body length is lacking from the formula. Results also suggest that BSA and the physiologic or pharmacologic factors that influence drug exposure may not be closely correlated.<sup>1,2</sup> To illustrate the problem with BSA dosing in dogs, a 30-kg (66-lb) dog has a BSA of 1 m<sup>2</sup>. If this dog were to be given doxorubicin at the recommended dosage of 30 mg/m<sup>2</sup>, this dog would be given a 30-mg total delivered dose, or 1 mg/kg. However, a 5-kg (11-lb) dog has a BSA of 0.30 m<sup>2</sup> and would be given a dosage of 9 mg (30 mg/m<sup>2</sup> times 0.30 m<sup>2</sup>) or 1.8 mg/kg (9 mg divided by 5 kg). This represents an 80% increase in dosage of

**Table 4. Equivalent surface area dosage conversion factors.<sup>11</sup>**

	To				
From	Mouse	Rat	Monkey	Dog	Man
Mouse	1	1/4	1/4	1/4	1/12
Rat	2	1	1/2	1/2	1/7
Monkey	4	2	1	3/5	1/3
Dog	6	4	5/3	1	1/2
Man	12	7	3	2	1

*Example: Given a dose of 50 mg/kg in the mouse, the appropriate dose in the monkey, assuming equivalency on the basis of mg/m<sup>2</sup>, is 50 mg/kg  $\times$  1/4 = 13 mg/kg.*

*This table gives approximate factors for converting doses expressed in terms of mg/kg from 1 species to an equivalent surface area dose expressed as mg/kg in the other species tabulated. The assumptions and constants of the formula by Freirich et al. (1966) are used.*



Table 5. Chemotherapy agents and dosing recommendations in exotics.

Drug	Dose	Tumor type
<b>Ferrets</b> <sup>21,22,26,31-37</sup>		
Vincristine*	0.75 mg/m <sup>2</sup> i.v.	Lymphoma
	2.0 mg/m <sup>2</sup> i.v.	Lymphoma
	0.12 mg/kg i.v.	Lymphoma
	0.20 mg/kg i.v.	Lymphoma
Cyclophosphamide†	200 mg/m <sup>2</sup> p.o., s.c.	Lymphoma
	10 mg/kg p.o.	Lymphoma
L-asparaginase‡	400 IU/kg s.c., i.m.	Lymphoma
Chlorambucil	1 mg/kg p.o.	Lymphoma
Doxorubicin*‡	20 mg/m <sup>2</sup> i.v.	Lymphoma, squamous cell carcinoma
	2 mg/kg i.v.	Lymphoma
Methotrexate	0.5 mg/kg i.v.	Lymphoma
Bleomycin	10 U/m <sup>2</sup> s.c.	Squamous cell carcinoma
<b>Avian</b> <sup>5,18,20,24,25,30,38-40</sup>		
Carboplatin	125 mg/m <sup>2</sup> i.v.	Bile duct carcinoma
	15 mg/kg i.o.	
Chlorambucil	1 mg/kg p.o.	Lymphocytic leukemia, hepatocellular carcinoma
Doxorubicin*‡	60 mg/m <sup>2</sup> i.v.	Osteosarcoma, hemangiosarcoma
Vincristine*	0.75 mg/m <sup>2</sup> i.v.	Lymphocytic leukemia
<b>Reptile</b> <sup>18</sup>		
Cytosine arabinoside	30 mg/kg s.c.	Lymphoma (may have caused severe toxicity)
<b>Rodent</b> <sup>18</sup>		
Doxorubicin (liposomal encapsulated)	6 mg/kg i.v.	Mammary adenocarcinoma
<b>Rabbit</b> <sup>41§</sup>		
Carboplatin	150-180 mg/m <sup>2</sup> i.v. q 3-4 weeks	Carcinoma
CCNU	50 mg/m <sup>2</sup> p.o. q 3-6 weeks	Lymphoma
Cyclophosphamide†	50 mg/m <sup>2</sup> p.o. q 24 hours for 2-3 days per week	Lymphoma
	100-200 mg/m <sup>2</sup> i.v. q 1-3 weeks (often combined with doxorubicin)	Lymphoma
Doxorubicin*‡	1 mg/kg i.v. q 2-3 weeks	Lymphoma
L-asparaginase‡	400 IU/kg i.m. or s.c.	Lymphoma
Mitoxantrone*	5-6 mg/m <sup>2</sup> i.v. q 3 weeks	Carcinoma
Prednisone	0.5-2.0 mg/kg p.o.	Lymphoma
Vincristine*	0.5-0.7 mg/m <sup>2</sup> i.v. q 1-2 weeks	Lymphoma

i.m. = intramuscularly; i.v. = intravenously; p.o. = by mouth; s.c. = subcutaneously; i.o. = intraosseous.

\*Drug must be administered intravenously via a clean stick to avoid extravasation and perivascular necrosis.

†Injectable cyclophosphamide can be administered orally at the same dose but may require dilution in propylene glycol for appropriate dosing. Alternately, an oral formulation can be compounded by a professional compounding pharmacy. This drug should be administered in the hospital to avoid unnecessary human contact or risk with the use of a liquid chemotherapeutic.

‡Premedicate with diphenhydramine, 1 to 2 mg/kg, 30 minutes before administration to prevent anaphylactic response.

§Personal observations.

doxorubicin when based on body weight. Doxorubicin clearly does not fit the BSA model, yet studies have not yet been completed to determine the

appropriate dosage based on body weight.<sup>3,4</sup> The conclusion, until such studies are conducted in animals of varying body size, shape, weight, and

length, is that chemotherapeutic drug dose selection in human and veterinary medicine remains anecdotal.

Thus, BSA is a difficult concept to define and is a variable that is extremely difficult to measure reproducibly. Several different formulae for predicting surface area in humans from measurements of height and weight have been derived (Table 1). The author prefer to use the modified Du Bois formula<sup>7</sup> (Table 2) for chemotherapy dosing in dogs and cats, and the Freireich formula<sup>11</sup> for scaling up from mouse and rat to other warm-blooded species or exotics (Tables 3 and 4). Despite anatomical, physiological, and biochemical differences among animal species, the pharmacokinetic disposition of many chemotherapy agents in avians,<sup>20</sup> reptiles,<sup>17</sup> ferrets,<sup>21,22</sup> and other exotics is similar in some respects to the kinetics reported previously in dogs<sup>3,6,16</sup> and humans.<sup>23</sup> Thus, it is likely that specific dosing requirements would be largely determined by the sensitivity of the tumor to the chemotherapy agent and by its toxicity, rather than any large-scale dosage alterations driven by significant pharmacokinetic differences. Current recommendations are to dose chemotherapy in avians as recommended in dogs<sup>3,6,16,20,24,25</sup> and to dose chemotherapy in ferrets as recommended for cats<sup>4,5,21,22,26</sup> (Table 5).

## References

1. Frazier DL, Price GS: Use of body surface area to calculate chemotherapeutic drug dose in dogs: II. Limitations imposed by pharmacokinetic factors. *J Vet Intern Med* 12(4):272-278, 1998
2. Price GS, Frazier DL: Use of body surface area (BSA)-based dosages to calculate chemotherapeutic drug dose in dogs: I. Potential problems with current BSA formulae. *J Vet Intern Med* 12(4):267-271, 1998
3. Arrington KA, Frazier DL, Tabeling GS, et al: The comparison of pharmacokinetics, hematologic parameters, and clinical signs of doxorubicin administration in small and large dogs, in: *Proceedings of the 12th Annual Conference of the Veterinary Cancer Society*, Asilomar, CA, 1992, pp 36-37
4. Hahn KA, Frazier DL, Cox SK, et al: Effect of infusion regime on doxorubicin pharmacokinetics in the cat. *J Am Anim Hosp* 33:427-433, 1997
5. Hahn KA, Nolan ML, McEntee MF, et al: Hematologic and systemic toxicoses associated with carboplatin administration in cats. *Am J Vet Res* 58:677-679, 1997
6. Bravo L, Hahn KA, Legendre AM, et al: Evaluation of pre- and post-cisplatin renal function in 23 tumor-bearing dogs, in: *Proceedings of the 12th Annual Conference of the Veterinary Cancer Society*, Asilomar, CA, 1992, pp 39-40
7. Du Bois D, Du Bois EF: A formula to estimate the approximate surface area if height and weight be known. *Arch Intern Med* 17:863-871, 1916
8. Baker SD, Verweij J, Rowinsky EK, et al: Role of body-surface area in dosing of investigational anticancer agents in adults: 1991-2001. *J Natl Cancer Inst* 94:1883-1888, 2002
9. Dooley MJ, Poole SG: Poor correlation between body surface area and glomerular filtration rate. *Cancer Chemother Pharmacol* 46:523-526, 2000
10. Gurney H: Dose calculation of anticancer drugs: A review of the current practice and introduction of an alternative. *J Clin Oncol* 14:2590-2611, 1996
11. Freireich EJ, Gehan EA, Rall DP, et al: Quantitative comparison of toxicity of anticancer agents in mouse, rat, hamster, dog, monkey, and man. *Cancer Chemother Rep* 50:219-244, 1966
12. Dedrick RL: Animal scale-up. *J Pharmacokinet Biopharm* 1:435-462, 1973
13. Grochow LB, Baraldi C, Noe D: Is dose normalization to weight or body surface area useful in adults? *J Natl Cancer Inst* 82:323-325, 1990
14. Reilly JJ, Workman P: Normalization of anti-cancer drug dosage using body weight and surface area: Is it worthwhile? *Cancer Chemother Pharmacol* 32:411-418, 1993
15. Sawyer M, Ratain MJ: Body surface area as a determinant of pharmacokinetics and drug dosing. *Invest New Drugs* 19:171-177, 2001
16. Ausmus P, Hahn KA, Avenell JS, et al: Pharmacokinetics of cisplatin (2 mg/kg v. 60 mg/m<sup>2</sup>) in 31 dogs, in: *Proceedings of the 14th Annual Conference of the Veterinary Cancer Society*, Townsend, TN, 1994, pp 43-44
17. Graham JE, Kent MS, Theon A: Current therapies in exotic animal oncology. *Vet Clin North Am Exotic Anim Pract* 7(3):757-782, 2004
18. Kent MS: The use of chemotherapy in exotic animals. *Vet Clin North Am Exotic Anim Pract* 7(3):807-820, 2004
19. Doolan M: Adriamycin chemotherapy in a blue-front Amazon with osteosarcoma, in: *Proceedings of the Annual Conference of the Association of Avian Veterinarians*, Reno, NV, 1994, pp 89-91
20. Filippich IJ, Bucher AM, Charles BG, et al: Intravenous cisplatin administration in sulphur-crested cockatoos (*Cacatua galerita*): Clinical and pathologic observations. *J Avian Med Surg* 15:23-30, 2001
21. Antinoff N, Hahn KA: Ferret oncology: Diseases, diagnostics, and therapeutics. *Vet Clin North Am Exotic Anim Pract* 7(3):579-626, 2004
22. Antinoff N: Neoplasia in ferrets, in Kirk RW, Bonagura JD (eds): *Current Veterinary Therapy XIII*. Philadelphia, PA, Saunders, 1999, pp 1149-1152
23. Gehan EA, George SL: Estimation of human body surface area from height and weight. *Cancer Chemother Rep* 54:225-235, 1970
24. Hahn KA, Jones MP, Petersen MG, et al: Clinical and pathological characterization of an osteoma in a barred owl. *Avian Pathol* 27:306-308, 1998
25. Hahn KA, Jones MP, Petersen MG, et al: Metastatic pheochromocytoma in a parakeet. *Avian Dis* 41:751-754, 1997
26. Hamilton TA, Morrison WB: Bleomycin chemotherapy for metastatic squamous cell carcinoma in a ferret. *J Am Vet Med Assoc* 198(1):107-108, 1991
27. Boyd E, Scammon RE, Lawrence D: The determina-

- tion of surface area in living children. *Proc Soc Exp Biol Med* 27:445-449, 1930
28. Mosteller RD: Simplified calculation of body-surface area. *N Engl J Med* 317:1098, 1987
  29. Haycock GB, Schwartz GJ, Wisotsky DH: Geometric method for measuring body surface area: A height-weight formula validated in infants, children and adults. *J Pediatr* 93:62-66, 1978
  30. France M, Gilson S: Chemotherapeutic treatment of lymphosarcoma in a Moluccan cockatoo, in: *Proceedings of the Annual Conference of the Association of Avian Veterinarians*, Nashville, TN, 1993, pp 15-19
  31. Brown SA: Neoplasia, in Hillyer EV, Quesenberry KE (eds): *Ferrets, Rabbits, and Rodents: Clinical Medicine and Surgery*. Philadelphia, PA, WB Saunders, 1997, pp 91-114
  32. Hutson CA, Kopit MJ, Walder EJ: Combination doxorubicin and orthovoltage radiation therapy, single agent doxorubicin, and high-dose vincristine for salvage therapy of ferret lymphosarcoma. *J Am Anim Hosp Assoc* 28(4):365-368, 1992
  33. Olsen GH, Turk MA, Foil CS: Disseminated cutaneous squamous cell carcinoma in a ferret. *J Am Vet Med Assoc* 186(7):702-703, 1985
  34. Rassnick KM, Gould WJ 3<sup>rd</sup>, Flanders JA: Use of a vascular access system for administration of chemotherapeutic agents to a ferret with lymphoma. *J Am Vet Med Assoc* 206(4):500-504, 1995
  35. Rodger BA: Possibility of virally induced lymphoma in ferrets. *Can Vet J* 24:237, 1980
  36. Rosenthal K: Ferrets. *Vet Clin North Am Small Animal Pract* 24(1):1-23, 1994.
  37. Williams BH, Weiss CA: Ferret Neoplasia, in Quesenberry KE, Carpenter JW, (eds): *Ferrets, Rabbits, and Rodents: Clinical Medicine* (ed 2). St. Louis, MO, Saunders, 2003, pp 91-106
  38. Freeman KP, Hahn KA, Jones MP, et al: Unusual presentation of an Amazon parrot (*Amazona amazona*) with hepatocellular carcinoma. *Avian Path* 28: 203-206, 1999
  39. Freeman KP, Hahn KA, Jones MP, et al: Right leg muscle atrophy and osteopenia caused by renal adenocarcinoma in a cockatiel (*Nympholopis undulatus*). *Vet Radiol Ultrasound* 40:144-147, 1999
  40. Manucy TK, Bennett RA, Greenacre CB, et al: Squamous cell carcinoma of the mandibular beak in a Buffon's macaw (*Ara ambigu*). *J Avian Med Surg* 12:158-166, 1998
  41. Heatley JJ, Smith AN: Spontaneous neoplasms of lagomorphs. *Vet Clin North Am Exotic Anim Pract* 7(3):561-578, 2004